

1 Interannual variability of the Subpolar Mode Water
2 properties over the Reykjanes Ridge during
3 1990-2006

V. Thierry,¹ E. de Boisséson,¹ H. Mercier²

V. Thierry, Ifremer, Laboratoire de Physique des Océans, UMR 6523 CNRS/Ifremer/UBO,
Ifremer Centre de Brest, BP 70, 29280 Plouzané cedex, France (vthierry@ifremer.fr)

¹Ifremer, Laboratoire de Physique des
Océans, UMR 6523 CNRS/Ifremer/UBO,
Brest, France

²CNRS, Laboratoire de Physique des
Océans, UMR 6523 CNRS/Ifremer/UBO,
Brest, France

4 **Abstract.** Combining hydrographic data from the OVIDE section (Greenland-
5 Portugal) with Argo and historical CTD data over the period 1990-2006, we
6 estimate the variability of the core properties of a variety of Subpolar Mode
7 Water (SPMW) observed on the eastern flank of the Reykjanes Ridge. This
8 SPMW acquires its core properties in the winter mixed layer along the east-
9 ern side of the Reykjanes Ridge. We find that the February SST along the
10 ridge is a proxy for its core temperature. The sources of this mode water are
11 water masses advected by the mean cyclonic circulation in the Iceland Basin.
12 A density compensated tendency for cooling and freshening of the SPMW
13 core properties is observed in the early 1990s. It stops in 1996 and is followed
14 by an increase in temperature and salinity (+ 1.41°C and + 0.11 psu) and
15 a decrease in density (- 0.12 kg m⁻³) until at least 2003. During the entire
16 period, the data do not show any significant modification in the depth of the
17 mode water core while they suggest that the thickness of the layer shrank.
18 The variability of the local air-sea freshwater and heat fluxes cannot explain
19 the observed salinity and temperature variations. They are most likely re-
20 lated to the modifications of the properties of the SPMW sources due to the
21 recently evidenced changes, driven by the North Atlantic Oscillation, in the
22 relative contributions of subtropical waters and subpolar waters in the Ice-
23 land Basin.

1. Introduction

24 Formed in deep winter mixed layers, mode waters are identified by nearly uniform prop-
25 erties in the vertical near the top of the permanent pycnocline. They cover large horizontal
26 areas in all oceans (see *Hanawa and Talley* [2001] for a review). Their locations and prop-
27 erties are set by complex interactions between air-sea fluxes of buoyancy and momentum,
28 circulation and mixing. The forcings being subject to interannual to interdecadal variabil-
29 ity, mode water characteristics vary at the same time scales. In the eastern North Atlantic,
30 *González-Pola et al.* [2005] show that Eastern North Atlantic Central Water (ENACW)
31 in the Bay of Biscay warmed and slightly freshened over the period 1992-2003 ($+0.032 \pm$
32 $0.008 \text{ }^\circ\text{C yr}^{-1}$ and $-0.002 \pm 0.001 \text{ psu yr}^{-1}$, respectively). The core density of this mode
33 water decreased from about 27.2 to 27.1 kg m^{-3} and the interannual salinity variations
34 seem to be related to the local P-E (precipitation minus evaporation) regime. Further
35 north, in the subpolar gyre of the North Atlantic, the process of transformation of the
36 warm, saline subtropical waters into intermediate and deep waters [*McCartney and Talley,*
37 *1982; Read, 2001; Perez-Brunius et al., 2004*] results in several varieties of Subpolar Mode
38 Water (hereafter SPMW) distributed around the gyre. SPMW along 20°W and north of
39 40°N are warmer ($\sim 0.7^\circ\text{C}$), saltier (~ 0.1) and lighter in 2003 than in 1993 [*Johnson and*
40 *Gruber, 2007*]. According to those authors, those changes are related to the NAO (North
41 Atlantic Oscillation), the dominant mode of atmospheric variability in the North Atlantic
42 sector. The densest variety of SPMW, the Labrador Sea Water (LSW), which is the main
43 contributor to the lower limb of the Meridional Overturning Circulation, is also subject to
44 large decadal property variations [*Dickson et al., 1996; Yashayaev, 2007*]. In the Labrador

45 Sea, the ocean heat loss decreased since the mid 1990s which limited the convection to the
46 upper 1200 m and led to the generation of a new salinity minimum layer and to a warming
47 and salinisation of the older deep LSW due to lateral mixing [*Bersch et al.*, 2007]. The
48 decadal variations of the convective activity in the Labrador Sea are correlated with the
49 NAO and because of the horizontal pattern of the atmospheric forcing, those variations
50 are anti-correlated with the convective activity in the Greenland Sea and in the Sargasso
51 Sea [*Dickson et al.*, 1996]. In this latter area, *Kwon and Riser* [2004] and *Peng et al.*
52 [2006] show that both temperature and subduction rate of the Subtropical Mode Water
53 are correlated with the NAO on decadal time scales, with NAO leading by 2-3 years.
54 Among the processes that account for the changes in the LSW properties, changes in the
55 properties of the water masses that enter the Labrador Sea are thought to be important.
56 Before any attempt to quantify the impact of the upstream water masses changes on the
57 LSW, we must document the properties and variability of those water masses, which is
58 the aim of this paper. We focus on a mode water found in the North Atlantic Ocean
59 over the Reykjanes Ridge (Fig. 1) because it lies in a central position along the path
60 of the subpolar gyre where exchanges between the eastern and western parts of the gyre
61 occur. It also contributes to the warm and salty waters that enter the Labrador Sea by
62 the West Greenland Current and that influence both convection and restratification in
63 the Labrador Sea [*Cuny et al.*, 2002; *Myers et al.*, 2007; *Yashayaev*, 2007]. In comple-
64 menting other studies on subpolar mode water variability that were undertaken either in
65 the eastern Atlantic [*Holliday*, 2003; *Johnson and Gruber*, 2007] or in the Labrador Sea
66 [*Dickson et al.*, 1996; *Yashayaev*, 2007] but neither in the central part of the subpolar
67 gyre, this work helps providing a basin scale view of the mode water variability in the

68 North Atlantic. Finally, documenting the variability of this mode water is also important
69 for models because it lies in a region where models have deficiencies in representing water
70 masses properties and circulation [*Treguier et al.*, 2005].

71 The NAO index is defined here as the principal component time series of the leading
72 EOF of winter (December through March) Sea Level Pressure anomalies over the Atlantic
73 sector (20-80°N, 90°W-40°E) [*Hurrell*, 1995]. The horizontal pattern of this index consists
74 of a north-south dipole with two centers of opposite sign located near Iceland and Azores,
75 respectively. The NAO index was in a high positive state at the beginning of the nineties.
76 It shifted to a negative value in the winter 1995/1996. Although it remained in a moderate
77 positive state over the period 1996-2006, the NAO index presented an overall downward
78 trend and occasionally reached negative values (Fig. 2a).

79 The NAO index variations are correlated to large-scale fluctuations in the air-sea fluxes
80 of heat, freshwater and momentum over the North Atlantic Ocean and to changes in
81 the ocean circulation (see *Visbeck et al.* [2003] for a review). Since the mid-nineties,
82 the winter mean momentum flux and winter mean heat loss averaged over the Iceland
83 Basin (52-63°N and 33-10°W) and the subpolar gyre (50-65°N and 45-15°W, not shown)
84 decreased (Fig 2b,c). These recent decadal changes in the air-sea fluxes induced a decrease
85 in the gyre intensity [*Flatau et al.*, 2003; *Häkkinen and Rhines*, 2004; *Hátún et al.*, 2005]
86 accompanied with a northwestward shift of the subarctic front in the central Iceland basin
87 (roughly identified in the 1990s by the position of the winter 7°C SST isotherm; see *Flatau*
88 *et al.* [2003] and Fig. 10 in Sec. 3.2) and with a modification of the relative contributions
89 in the Iceland Basin of cold and low-saline waters of subpolar origin and warm and salty
90 waters of subtropical origin [*Bersch*, 2002; *Hátún et al.*, 2005]. According to *Johnson and*

91 *Gruber* [2007], these latter changes mainly explain the mode water variability along 20°W.
92 On these decadal time scales, the ocean response to the NAO is complex with significant
93 changes near strong mean current systems [*Visbeck et al.*, 2003]. On interannual time
94 scales, however, the ocean variability is dominated by NAO-induced changes in the air-
95 sea fluxes [*Visbeck et al.*, 2003].

96 The OVIDE project (Observatoire de la Variabilité Interannuelle à Decen-
97 nale/Observatory of the Interannual to Decadal Variability) repeats a trans-oceanic hy-
98 drographic section across the North Atlantic every other year since 2002 in order to
99 monitor and understand the low-frequency fluctuations of the oceanic Atlantic Merid-
100 ional Overturning Cell, heat and tracer transports and water mass characteristics in the
101 North Atlantic Ocean [*Lherminier et al.*, 2007]. The OVIDE section consists of full-water
102 column hydrographic stations between Portugal and the southern tip of Greenland (Cape
103 Farewell) (Fig. 1). The western part of the section is coincident with the A01E section
104 repeated several times since 1991 between Cape Farewell and Ireland (Fig. 1 and Ta-
105 ble 1). The common part of the two sections samples the Irminger Basin and part of
106 the Iceland Basin. It crosses the Reykjanes Ridge around 59°N where a thick pycnostad,
107 highlighting the presence of a SPMW variety, is clearly present near 300-500 m depth (Fig.
108 3). The aim of this paper is to investigate the interannual variability of this mode water
109 over the period 1990-2006. This is made possible because an adequate time series has
110 been created in combining CTD (conductivity-temperature-depth) measurements from
111 hydrographic cruises (OVIDE, A01E and few others, see Sec. 2) and Argo profiling floats.
112 Owing to the Argo array, we can also document properties of the SPMW in the northern

113 North Atlantic (50-66°N, 45-0°W) over the Argo period 2001-2006 to put the mode water
114 observed on the Reykjanes Ridge in a wider spatial context.

115 The data set is presented in Section 2. The mode water over the Reykjanes Ridge is
116 described in section 3. We also describe the interannual variability of its properties and
117 we discuss the source and formation area of this mode water. Section 4 discusses the
118 results and we conclude in Section 5.

2. Dataset and Mode Water Identification

119 The high-quality hydrographic stations carried on during the OVIDE 2002, 2004 and
120 2006 sections (Table 1) used a Neil Brown Mark III CTD02 probe. The rosette was
121 equipped with 28 8-liter bottles for tracers measurements and calibration purpose. The
122 CTD02 measurement accuracies are thought to be better than 1db for pressure, 0.002°C
123 for temperature, 0.003 for salinity and $1\mu\text{mol kg}^{-1}$ for dissolved oxygen [*Branellec et al.*,
124 2004]. High-quality CTD data from the A01E WOCE section (which is also referred to
125 as AR07E section) collected in 1991, 1992, 1995, 1996 and 1997 [*Bersch*, 1995; *Bersch*
126 *et al.*, 1999; *Bersch*, 2002; *Bersch et al.*, 2007] complement the OVIDE data, as well as
127 data from two additional cruises realized in 1990 as part of the NANSEN project [*van*
128 *Aken and Becker*, 1996] and in 1991 during the CONVEX-91 survey [*Read*, 2001] (Fig. 1
129 and Table 1).

130 Argo data downloaded from the Coriolis data center (<http://www.ifremer.fr/coriolis/>)
131 complement the dataset. The Coriolis data center provides quality-controlled in-situ data
132 in real-time and delayed mode and is a gateway to the global Argo data. Three levels
133 of quality control are performed to the Argo data. First, a series of standard automatic
134 quality control (QC) is applied (see the Argo quality control manual [*Argo*, 2005] for

135 more details). As the automatic real-time quality control procedure cannot identify small
136 salinity drifts or offsets that Argo floats experience due in particular to biofouling [*Wong*
137 *et al.*, 2003; *Boehme and Send*, 2005], a second quality control is performed at Coriolis,
138 following *Gaillard et al.* [2007], to remove dubious profiles. Finally, the delayed-mode pro-
139 cedure [*Wong et al.*, 2003; *Boehme and Send*, 2005] is applied to correct (when necessary)
140 offsets and drifts and to generate a qualified Argo dataset [*Argo*, 2005]. Among the 4578
141 Argo profiles downloaded for this analysis, half contain delayed-mode salinity data and
142 25% (about 600 profiles) have been corrected. Since the beginning of the Argo program,
143 float and sensor technology has been improved and it is expected that this percentage
144 will decrease with the replacement of the old fleet by new generation of floats. At the
145 time of our analysis, Argo profiles from SOLO floats with FSI CTD may have incorrect
146 pressure values [*Schiermeier*, 2007]. They have been excluded from our dataset. Real
147 time and delayed-mode Argo data containing both temperature (T) and salinity (S) are
148 considered in this study. T and S have a nominal accuracy of 0.01°C and 0.01 psu. In
149 case of duplicates, delayed-mode profiles replace real time data.

150 We will show in this study that the data from the Argo/Coriolis database provide
151 results that are fully consistent with that deduced from the ship-based high-quality CTD
152 measurements (see Sec. 3.1 and Fig. 7). This gives good confidence for using this dataset
153 in the future for the monitoring of the mode water properties.

154 During summer, mode waters are isolated from the atmosphere and their properties do
155 not evolve much, which allows the robust characterization of their properties. They are
156 characterized by a thick pycnostad between the seasonal and the permanent pycnocline
157 and can be identified by a minimum in the potential vorticity q [*Hanawa and Talley*,

2001]. For each profile collected from June through September (Fig. 1), the mode waters
are identified as the layer where $q < 6 \times 10^{-11} \text{ m}^{-1} \text{ s}^{-1}$ [Johnson and Gruber, 2007].
Analyzing the WOCE dataset collected in 1997, Talley [1999] uses a different criterion
($q < 4 \times 10^{-11} \text{ m}^{-1} \text{ s}^{-1}$) but that of Johnson and Gruber [2007] appears more adequate
for the identification of recent mode water vintages that are more stratified than those
formed at the beginning of the 90s (Fig. 5). The LSW is excluded in considering only
the layers where the potential density is less than 27.7 kg m^{-3} . A visual inspection is
then performed to eliminate the selected profiles that do not contain SPMW. Finally, we
retain the thickest layers in imposing that the thickness of the mode water layer must be
greater than 100 db. For each profile and in the layer satisfying the above conditions, the
core properties of the mode water are defined at the level where the potential vorticity is
minimum. Examples are displayed on Fig. 5. Potential temperature (θ), potential density
(σ_0) and q are deduced from the T and S profiles. θ and σ_0 are referenced to 0 db. A 4th-
order Butterworth filter with a cut-off wave length of 50 dbar is applied to the potential
vorticity estimated from the 1-db vertical resolution ship-based CTD measurements. Due
to the coarser vertical sampling, the Argo data are not filtered.

3. The Subpolar Mode Water over the Reykjanes Ridge

3.1. Properties and Interannual Variability

The profiles collected from 2001 to 2006 during the Argo period allow us to determine
the localization and the properties of the SPMW in the northern North-Atlantic (Fig.
6). SPMW are distributed around the subpolar gyre. They are found south of Rockall
Plateau and in Rockall Trough, in the northern part of the Iceland Basin, on the eastern
flank of the Reykjanes Ridge and in the northern part of the Irminger Basin. This picture

179 is fully consistent with an analysis based on data collected in 1997 by *Talley* [1999]. In the
180 eastern Iceland Basin, the density (salinity and potential temperature) of SPMW increases
181 (decrease) northward from 27-27.1 kg m⁻³ (35.5-35.6 and 11-12°C) at the southern limit
182 of the Rockall Trough to 27.4-27.5 kg m⁻³ (35.1-35.2 and 7-8°C) in the northern Iceland
183 Basin. The SPMW are absent in the central part of this basin. There, well-stratified
184 water masses are embedded within one of the three main branches of the North Atlantic
185 Current (NAC) [*Talley*, 1999; *Read*, 2001]. Along the Reykjanes Ridge, the mode water
186 variety has the following properties: $\sigma_0 \sim 27.4-27.5 \text{ kg m}^{-3}$, $\theta \sim 7-8^\circ\text{C}$ and $S \sim 35.1-35.2$.
187 The densest variety (excluding LSW) is observed in the northern part of the Irminger
188 basin with a density greater than 27.5 kg m⁻³ and salinity and potential temperature
189 usually lower than 35.1 and 7 °C, respectively.

190 Let us now consider the SPMW observed over the Reykjanes Ridge (it is called Atlantic
191 Water by *Read* [2001]). A comparison of the properties of this SPMW with previous
192 estimates shows large variability. The pool of uniform salinity water associated with this
193 SPMW was more saline by about 0.1 in 2004 than in 1992 (Fig. 4). Also, with a density
194 greater than 27.5 kg m⁻³, this SPMW was denser in 1992 (Fig. 3b) and in 1997 [*Talley*,
195 1999] than over the period 2001-2006 when the density was less than 27.5 kg m⁻³ (Figs.
196 3a, 5 and 6).

197 Our data set allows us to investigate the interannual variability of the properties of the
198 SPMW observed over the Reykjanes Ridge in a box located on the eastern flank of this
199 ridge (57.5-59.5°N, 31.5-28.5°W) (Fig. 1). For this purpose, we define the yearly property
200 value as the median of all estimates of the mode water properties in the box for a given
201 year (from summer profiles). We use the median instead of the mean because we do not

202 want to bias the result toward extreme values. The time-averaged of the yearly properties
203 between 1990 and 2006 are $\sigma_0 = 27.51 \text{ kg m}^{-3}$, $\theta = 7.07^\circ\text{C}$ and $S=35.13$. The core of this
204 mode water is located at 450 db. The time evolution of the core properties of this SPMW
205 is depicted in Fig. 7. Removing the corresponding yearly values from each estimates, the
206 standard deviation over 1990-2006 is estimated to 0.02 kg m^{-3} for σ_0 , 0.2°C for θ , 0.02 for
207 S and 73 db for the pressure at the mode water core. As not enough data are available to
208 provide an accurate confidence interval for each yearly estimate, we consider that changes
209 in the yearly values of the mode water properties are significant when they are greater
210 than 2 times the estimated standard deviation.

211 From 1990 to 1995, mode water properties are relatively stable ($\sigma_0 = 27.56 \text{ kg m}^{-3}$),
212 although a slight density compensated trend toward fresher and colder mode water is
213 observed (by about 0.04 and 0.24°C) (Fig. 7). The trend reversed in 1996. From 1996 to
214 2003, the salinity and the temperature of the mode water core increased. Those changes
215 are not density compensated and during the same time the density decreased. In 2003,
216 the SPMW was 1.41°C warmer, 0.11 saltier and 0.12 kg m^{-3} lighter than in 1995. The
217 warming and salinisation ceased after 2003. The data even suggest that the trend has
218 reversed since that year, but this has to be taken with caution as the 2003 yearly values
219 are deduced from two profiles only. However, this is in full agreement with measurements
220 collected along the AR7E section in 2003 and 2005 that shows the SPMW at 500 m depth
221 over the Reykjanes Ridge was cooler and fresher in 2005 than in 2003 [ICES, 2006].

222 A gap in our data set does not allow us to describe the time evolution of the properties of
223 this mode water between 1997 and 2002. It can be indirectly documented in considering
224 the time series of the February Reynolds SST [Reynolds *et al.*, 2002] averaged in the

225 Reykjanes Box that follows fairly well the SPMW core temperature (Fig. 7, see also Sec.
226 3.2 for more details) and measurements collected in 1999 along the A01E section. Those
227 data are not available for our analysis but are discussed by *Bersch* [2002] and *Bersch*
228 *et al.* [2007]. The February Reynolds SST exhibits a warm anomaly in 1998-1999 and a
229 visual inspection of the salty anomaly in the upper layers of the A01E section near 30°W
230 (see Fig. 10 in *Bersch et al.* [2007]) reveals the presence of a positive anomaly in 1999.
231 The long term trend in the core properties of the SPMW observed over the Reykjanes
232 Ridge are thus modulated by interannual variability, with warm and salty anomalies in
233 1998-1999.

234 According to our dataset, no significant change in the depth of the mode water core
235 occurred over the period 1990-2006, except in 2004 when the core was anomalously shallow
236 (Fig. 7). In the Bay of Biscay, the ENACW core remains also at the same depth over 1992-
237 2003 [*González-Pola et al.*, 2005] because, simultaneously, the isopycnal levels deepened
238 and the core density of this mode water decreased from about 27.2 to 27.1 kg m⁻³. We
239 expect that the same process explains the stability of the SPMW core depth over 1990-
240 2006. Finally, the data suggest that the thickness of the mode water layer was greater
241 than 300 db before 1996, while it has been usually lower than this value since then (Fig.
242 8). The extreme value observed in 2002 (thickness ~ 800 db) is due to an eddy sampled
243 during the 2002 OVIDE section. The mode water in this eddy presents extreme properties
244 with a density and a potential temperature at the core of the mode water less than 27.4
245 kg m⁻³ and greater than 8°C (Fig. 7). The two profiles displayed on Fig. 5 illustrate
246 fairly well the contrast in SPMW properties between the recent years and the beginning
247 of the 1990s.

3.2. Source and Formation Area

248 The Hydrobase 2 Atlas shows that the climatological 27.45, 27.5 and 27.55 kg m⁻³
249 isopycnals outcrop in winter southwest of Iceland parallel to the Reykjanes Ridge between
250 the 1000 and 2000 m isobaths (Fig. 9), which identifies the potential formation region of
251 the SPMW observed over the Reykjanes Ridge. Considering two boxes located along
252 the outcropping region of the 27.45-27.55 kg m⁻³ isopycnals (Fig. 9), we show that the
253 February SST averaged in these boxes and their time evolution are in fair agreement with
254 those in the Reykjanes box (Fig. 7). This was expected since the SST isotherms are
255 also parallel to the ridge (Fig. 10), and confirms uniform surface properties along the
256 ridge. In addition, the February SST interannual variability along the Reykjanes Ridge
257 follows the interannual variations of the SPMW core temperature in the Reykjanes box
258 (Fig. 7). Since the February SST over the eastern flank of the Reykjanes Ridge, which
259 represents the late winter mixed layer temperature, is also a proxy for the SPMW core
260 temperature, we conclude that the SPMW is formed in the winter mixed layer over the
261 eastern Reykjanes Ridge. Surface isotherms and isopycnals being parallel to the ridge, we
262 expect uniform mode water properties between southwest of Iceland and the Reykjanes
263 box, which is evidenced from the in situ data (Fig. 6).

264 In the Iceland Basin, the upper ocean waters overlying LSW are a mixture of cold, fresh
265 subarctic water masses and warm, saline subtropical water masses. From a water mass
266 point of view, there is some evidence that the warm and salty subtropical waters spread
267 northward in the northeastern North Atlantic [*Bower et al.*, 2002], circulate around the
268 northern Iceland Basin and flow southwestward along the eastern flank of the Reykjanes
269 Ridge [*Read*, 2001; *Pollard et al.*, 2004]. This is confirmed by *Bower et al.* [2002] who, using

270 acoustically tracked floats, estimated the mean circulation in the subpolar North Atlantic
271 on the 27.5 kg m^{-3} isopycnal (level of the mode water core) over 1993-2001 (Fig. 11).
272 After crossing the Mid-Atlantic Ridge between 50°N and 53°N , the NAC turns northward
273 in the Iceland Basin and then splits into two main branches. One branch turns sharply
274 anticlockwise to feed directly the Irminger Current on the western side of the Reykjanes
275 Ridge while the other branch, after penetrating farther north into the Iceland Basin,
276 returns southwestward along the eastern flank of the ridge and eventually crosses the
277 Reykjanes Ridge to feed the Irminger Current. Although, the surface circulation pattern
278 deduced from surface drifters drogued at 15 m depth exhibits an undefined mean flow
279 along the eastern flank of the Reykjanes Ridge [*Reverdin et al.*, 2003; *Flatau et al.*, 2003],
280 some of the 15-m drogued floats that were deployed south of Iceland moved southwestward
281 along the eastern flank of the ridge [*Reverdin et al.*, 2003]. This near-surface circulation
282 along the ridge would be better defined in winter (when the 27.5 kg m^{-3} isopycnal reaches
283 the surface) than in summer which corroborates a southwestward circulation along the
284 eastern flank of the Reykjanes Ridge on the 27.5 kg m^{-3} isopycnal. There is thus evidence
285 from water masses and circulation that the SPMW observed over the Reykjanes Ridge
286 is at least partly supplied by waters advected by the mean circulation from the northern
287 and eastern Iceland Basin.

288 This latter conclusion does not mean that this SPMW is directly connected to the
289 lighter SPMW variety lying in the eastern side of the Iceland Basin as hypothesized by
290 [*McCartney and Talley*, 1982]. *Brambilla* [2007] shows that the connection between the
291 (lighter) SPMW in the eastern Iceland basin and the (denser) SPMW over the Reykjanes
292 Ridge is unlikely, although it might occur intermittently. *Read* [2001] also concludes

293 that the SPMW properties are "set primarily by modification of whatever segment of the
294 temperature/salinity curve has reached the surface rather than by cooling and freshening
295 of central water along an advective path as described by McCartney and Talley [1982]".
296 Beside deep winter mixing and advection, the other factors that potentially help setting
297 the SPMW properties are mixing with underlying and lateral water masses and eddy
298 activity [Read, 2001].

4. Discussion

299 The warming and salinisation trend of the SPMW over the Reykjanes Ridge started in
300 1996 after the abrupt drop of the NAO index and persisted until at least 2003. During
301 that period, the NAO index presented a decreasing trend and the air-sea forcing changed
302 accordingly. We thus investigate the relationship between the properties changes of this
303 SPMW and the NAO-driven circulation and atmospheric forcing field variability.

304 Changes in local air-sea forcings are quantified in the formation area of this mode water
305 by averaging atmospheric fluxes in a box covering the northern part of the Iceland basin
306 (52-63°N, 33-10°W, Fig. 2). We first consider the freshwater flux because its relation
307 with the mixed layer salinity is straightforward due to the absence of feedback between
308 surface salinity and evaporation or precipitation. The annual P-E anomaly exhibits a
309 positive trend over the period 1990-2006 (Fig. 2), which excludes P-E as the driver of the
310 salinisation of the SPMW over the period 1995-2003. This result is in full agreement with
311 *Hátún et al.* [2005] and *Holliday* [2003], but it differs from *Josey and Marsh* [2005]' and
312 *González-Pola et al.* [2005]' findings. *Josey and Marsh* [2005] show that on an interdecadal
313 time scale, the freshening (~ 0.2) of the surface layers of the eastern half of the North
314 Atlantic subpolar gyre from the mid 1970s until the 1990s can be largely explained by an

315 increase in P-E in the gyre region. In the Bay of Biscay, the P-E regime on an interannual
 316 time scale seems to be the main driver for the ENACW salinity variations (in the range of
 317 0.05 to 0.1) over 1992-2003 [*González-Pola et al.*, 2005]. Understanding those differences
 318 on the role of P-E (both geographically and at different temporal scales) deserves to be
 319 investigated in detail but is beyond the scope of this study.

320 Let us now provide some insight on the possible effect of the local air-sea fluxes vari-
 321 ability on the winter SST and on mode water temperature. The NAO index was positive
 322 beginning of the 1990's and the winter heat loss and the winter zonal momentum were
 323 maximum during that period. The NAO index decrease since the mid-1990s is accompa-
 324 nied in the subpolar gyre with a decrease in both the zonal momentum and the total heat
 325 loss and an increase in the SST as revealed by the westward shift of the position of the
 326 February 7°C isotherm since 1996 (Fig. 10). In order to quantify the ocean response to
 327 the air-sea heat flux variability, we compute, for each profile in the Reyjkanes Box, the
 328 heat content anomaly relative to a reference temperature in the mode water layer as :

$$H = \int \rho C_p (T - T_R) dz \quad (1)$$

329 with ρ and C_p the density and the specific heat capacity of seawater, z the depth in
 330 metres, T the potential temperature and T_R the reference temperature which is chosen as
 331 the mean temperature of the mode water core over 1990-2006 (here 7.07°C). The mode
 332 water layer is deduced from the mode water thickness and the depth of the mode water
 333 core and varies from one profile to the other. A yearly value is defined as the median of
 334 all estimates of the heat content anomaly in the box for a given year. We then compare
 335 the annual variations of this heat content anomaly to changes in the annual air-sea heat

336 flux multiplied by time following *Holliday* [2003]. With variations of order 0.5 J m^{-2} over
337 the period 1995-2006 compared to more than 2 J m^{-2} for the heat content anomaly, we
338 estimate that the local heat flux variability is a minor contribution to the SPMW core
339 temperature variations (Fig. 12).

340 Long-term changes in water mass properties have been reported in the whole eastern
341 subpolar gyre over the last decade. *Hátún et al.* [2005] show clearly a continuous salinisa-
342 tion of the Atlantic Inflow to the Nordic Seas over 1995-2004 by about 0.1 psu. Analyzing
343 SPMW property variations along 20°W , *Johnson and Gruber* [2007] observe extreme and
344 opposite conditions in 1993 (colder and less saline) and in 2003 (warmer and saltier),
345 while intermediate conditions are observed in 1988 and 1998. Analyzing data from the
346 Extended Ellet Line in the northern Rockall Trough over the period 1975-2000, *Holliday*
347 [2003] shows that temperature and salinity exhibit coherent decadal fluctuations with
348 highs in the mid 1980s and late 1990s and lows in the late 1970s and early 1990s. Recent
349 measurements along the Extended Ellet Line in the Rockall Trough show that since the
350 late 1990s the temperature and salinity continue to increase (see these unpublished data
351 on <http://www.noc.soton.ac.uk/obe/PROJECTS/EEL>, hereafter EEL web site): over
352 the period 1995-2005, the changes are of order 0.8°C and 0.08 psu. The air-sea heat and
353 freshwater fluxes cannot explain the amplitude of the temperature and salinity variations
354 [*Holliday*, 2003; *Hátún et al.*, 2005] or the deep penetration of the changes [*Johnson and*
355 *Gruber*, 2007]. In all cases, the authors conclude that the water mass variability is primar-
356 ily attributable to NAO-related changes in the shape and strength of the subpolar gyre
357 and in the regional circulation that modify the relative contribution of relatively fresh
358 and cold water masses of subpolar origin (SubArctic Intermediate Waters) and warmer

359 and saltier water mass of southern origin (Western North Atlantic Central Water that
360 are transported by the North Atlantic Current and Eastern North Atlantic Central Water
361 that comes from the intergyre area in the eastern North-Atlantic). *Häkkinen and Rhines*
362 [2004] suggest that the gyre weakening after 1995 is primarily attributable to changes in
363 the net heat flux. Using models, *Böning et al.* [2006] suggest that the wind stress is also
364 a contributor to the gyre variability especially in the early 1990s when both the net heat
365 flux and the wind stress acted in concert to produce an intense transport.

366 The variability of the SPMW observed over the Reykjanes Ridge is fully consistent with
367 the variability of the Atlantic Inflow to the Nordic Seas [*Holliday, 2003; Hátún et al., 2005*]
368 and the variability of the mode water south of Iceland along 20°W [*Johnson and Gruber,*
369 2007]. All water masses exhibit coherent temperature and salinity variations and tend to
370 be lighter, saltier and warmer since 1996. The variations are of order 0.1 in salinity and
371 1°C in 10 years. According to the mean circulation pattern on the 27.5 kg m⁻³ isopycnal
372 (Fig. 11), to the fact that the SPMW over the Reykjanes Ridge is supplied by waters from
373 the northern and eastern Iceland Basin and that the local P-E and heat flux variations
374 in the formation area of this SPMW cannot explain the temperature and salinity changes
375 over 1990-2006, we conclude that the variations of the core properties of this SPMW are
376 likely mainly due to the variations in the properties of its source water masses.

377 The long-term trend in the core properties of the SPMW observed over the Reykjanes
378 Ridge is modulated by interannual fluctuations. Warm and salty anomalies are observed
379 in 1998-1999 and possibly in 2003 (Sec. 3.1). A peak in salinity is also observed in the
380 upper layers of the Rockall Trough in 1998 and in 2003 (*Hátún et al.* [2005], see also the
381 EEL web site). Those anomalies are lagged by one year with the anomalies reported by

382 *Bersch et al.* [2007] in the central part of the Iceland Basin in 1996-1997 and in 2002.
383 According to these authors, the fast response of the Subarctic Front in the Iceland Basin
384 to the drop of the NAO index during the winters 1995/1996 and 2001/2002 (Fig. 2a)
385 induces a northwestward shift of the front with a time lag of 1 to 2 years. In 1999, when
386 the NAO index returned to a positive value, the low saline subarctic water masses began
387 to occupy again the Iceland Basin (see Fig. 10 in *Bersch et al.* [2007]). The interannual
388 anomalies observed on the eastern flank of the Reykjanes Ridge, lagged to the NAO index
389 by 2-3 years, are likely linked to those reported by *Bersch et al.* [2007].

390 The SPMW core warmed by about 1.4°C in 9 years which is one order of magnitude
391 greater than the temperature increase of 0.274°C in the top 700 m depth of the North-
392 Atlantic over 1955-2003 reported by *Levitus et al.* [2005]. Part of the observed changes
393 might be related to this global warming but yet, the length of our time series and the
394 volume of water considered here do not allow us to separate long term changes due to
395 anthropogenic influence from intrinsic oceanic variability. For this purpose, we believe it
396 is worth continuing the monitoring of the SPMW core properties in the North Atlantic in
397 using ship-based hydrographic data and Argo data.

5. Conclusion

398 Combining CTD data from different hydrographic sections and Argo data collected in
399 the northern North-Atlantic over 1990-2006, we provide a picture of the geographical
400 distribution of the subpolar mode waters in the North Atlantic. The core property of the
401 subpolar mode waters is individually identified in profiles collected from June through
402 September. In particular, a variety of mode water is identified along the eastern flank
403 of the Reykjanes Ridge. Its mean properties over 1990-2006 are $\sigma_0 = 27.51 \text{ kg m}^{-3}$, $\theta =$

404 7.07°C and $S=35.13$. According to water mass properties and to the mean circulation
405 pattern on the 27.5 kg m⁻³ isopycnal, we conclude that the sources of this SPMW are
406 advected by the mean circulation from the northern and eastern Iceland Basin and that
407 this SPMW acquires its final properties in the winter mixed layer southwest of Iceland on
408 the eastern side of the Reykjanes Ridge roughly between the 1000 and 2000 m isobaths.
409 We also show that the February SST along the Reykjanes Ridge is a proxy for the SPMW
410 core temperature.

411 Our data set allows us to investigate the interannual variability of this SPMW in a
412 box centered near 58.5°W and 30°W. The density compensated tendency for cooling and
413 freshening observed in the early 1990s is interrupted in 1996 when the trend reversed until
414 at least 2003. During that period, this SPMW warmed by 1.41°C and became more saline
415 by 0.11. As a consequence, the density of the SPMW core decreased by 0.12 kg m⁻³. Since
416 2003, the properties of this mode water are relatively stable with possibly a slight trend
417 toward colder and fresher properties. In combining February SST data and results from
418 data collected in 1999 along the A01E section, we suggest that the core properties of this
419 mode water reached a local maximum (warm and salty anomaly) in 1998-1999. During
420 the whole period (1990-2006), the data do not show any significant modifications in the
421 depth of the mode water core but they suggest that the thickness of the layer shrank.

422 The warming and salinisation that are observed after 1995 occurred simultaneously
423 with large changes in the NAO index that was largely positive until 1995 and, after a
424 large negative value in 1996, never returned to large positive values. During the same
425 time, the winter air-sea heat fluxes and momentum fluxes in the northern Iceland Basin
426 decreased leading to decreased winter heat loss, warmer SST and potentially warmer

427 SPMW. However, we show that the local variations in the air-sea fluxes are a minor
428 contribution to the warming trend. In addition, the annual freshwater flux exhibits a
429 positive trend over 1990-2006 and cannot explain the observed salinity changes. The
430 decrease in NAO index is associated with an invasion of warm and salty waters in the
431 upper layers of the eastern subpolar gyre and in the Iceland Basin. Since the simultaneous
432 changes in temperature and salinity cannot be explained by variations in local air-sea
433 fluxes, we conclude that they are most likely due to the displacements of water masses
434 associated with changes in gyre circulation and shape.

435 The long term trend of the core properties of the SPMW observed over the Reykjanes
436 Ridge is modulated by interannual variations: warm and salty anomalies are observed in
437 1998-1999 and possibly in 2003. Those anomalies could be related, with a time lag of 2-3
438 years, to the abrupt drop of the NAO index that occurred in 1995/1996 and 2000/2001
439 and that induced a northwestward retreat of the Subarctic Front in the Iceland basin 1-2
440 years after the shift.

441 Anthropogenic variability (global warming) surperimposes to the intrinsic ocean vari-
442 ability but longer time series are needed to disentangle human-driven long-term trend from
443 the natural oscillation of the system. This is an important issue as water mass properties
444 variations in the subpolar gyre can have a large impact on the distribution and habitat
445 of some fish species for instance [*Pedchenko*, 2005]. In the context of climate change,
446 we are thus looking for some indicator of the oceanic and atmospheric state and mode
447 water properties in the North-Atlantic ocean are a good candidate [*Banks and Wood*,
448 2002]. Indeed, they contribute to the preconditionning of the water masses before deep
449 convection in the Nordic, Irminger and Labrador Seas, they contribute to the heat and

450 CO₂ storage in the ocean and they can be considered as an integrator of the oceanic and
451 atmospheric variability. The monitoring of this mode water properties will be continued
452 in the future owing to perennial observations (OVIDE project, Argo floats) and the role
453 of each mechanisms presented in this paper will be investigated in details in using both
454 data and models.

455 **Acknowledgments.** Virginie Thierry is supported by IFREMER (Institut Français
456 de Recherche pour l'Exploitation de le Mer), Eric de Boisséson is funded by IFRE-
457 MER and Météo France and Herlé Mercier is funded by CNRS (the French Centre Na-
458 tional de la Recherche Scientifique). Support to the OVIDE project comes from IFRE-
459 MER, CNRS and INSU (Institut National des Sciences de l'Univers) and from French
460 national programs (GMMC and LEFE-IDAO). OVIDE is a contribution to CLIVAR.
461 NCEP Reanalysis data are provided by the NOAA-CIRES Climate Diagnostics Cen-
462 ter, Boulder, Colorado, USA, from their Web site at <http://www.cdc.noaa.gov/>. NAO
463 Index data are provided by the Climate Analysis Section, NCAR, Boulder, USA and
464 downloaded from <http://www.cgd.ucar.edu/cas/jhurrell/indices.data.html#naopcdjfm>.
465 Hydrobase 2 are data provided by Ruth Curry (WHOI) and downloaded from
466 http://www.whoi.edu/science/PO/hydrobase/HB2_home.htm. The OI.v2 monthly
467 Reynolds SST data are provided by the NOAA and downloaded from <http://www.emc.ncep.noaa.gov/research/>

References

468 Argo (2005), Argo quality control manual V 2.1 - ar-um-04-01.
469 Banks, H., and R. Wood (2002), Where to look for anthropogenic climate change in the
470 ocean, *15*, 879–891.

- 471 Bersch, M. (1995), On the circulation of the northeastern North Atlantic, *Deep Sea Res.*
472 *I*, *42*, 1583–1607.
- 473 Bersch, M. (2002), North Atlantic Oscillation-induced changes of the upper layer cir-
474 culation in the northern North Atlantic Ocean, *J. Geophys. Res.*, *107*, 3156, doi:
475 10.1029/2001JC000901.
- 476 Bersch, M., J. Meincke, and A. Sy (1999), Interannual thermohaline changes in the north-
477 ern North Atlantic 1991-1996, *Deep Sea Res. II*, *46*, 55–75.
- 478 Bersch, M., I. Yashayaev, and K. P. Koltermann (2007), Recent changes of the ther-
479 mohaline circulation in the subpolar North Atlantic, *Ocean Dyn.*, *57*, 223–235, doi:
480 10.1007/s10236-007-0104-7.
- 481 Boehme, L., and U. Send (2005), Objective analyses of hydrographic data for referencing
482 profiling float salinities in highly variable environments, *Deep Sea Res. II*, *52*, 651–
483 664, doi:10.1016/j.dsr2.2004.12.014.
- 484 Böning, C. W., M. Scheinert, J. Dengg, A. Biastoch, and A. Funk (2006), Decadal variabil-
485 ity of subpolar gyre transport and its reverberation in the North Atlantic overturning,
486 *Geophys. Res. Lett.*, *33*, L21S01, doi:10.1029/2006GL026906.
- 487 Bower, A. S., B. L. Cann, T. Rossby, W. Zenk, J. Gould, K. Speer, P. L. Richardson,
488 M. D. Prater, and H.-M. Zhang (2002), Directly measured mid-depth circulation in the
489 northeastern North Atlantic Ocean, *Nature*, *419*, 603–607.
- 490 Brambilla, E. (2007), The upper limb of the North Atlantic overturning circulation: in-
491 vestigation of the subtropical-subpolar gyre exchange and Subpolar Mode Water, Uni-
492 versity of California, San Diego, 148pp.

- 493 Branellec, P., A. billant, and H. Mercier (2004), Campagne OVIDE 2002-Rapport de
494 données CTD-O2, Rapport interne DOPS/LPO/04-01.
- 495 Cuny, J., P. B. Rhines, P. P. Niiler, and S. Bacon (2002), Labrador Sea boundary currents
496 and the fate of Irminger Sea Water, *J. Phys. Oceanogr.*, *32*, 627–647.
- 497 Dickson, R., J. Lazier, J. Meincke, P. Rhines, and J. Swift (1996), Long term coordinated
498 changes in the convective activity of the North Atlantic, *Prog. Ocean.*, *38*, 241–295.
- 499 Flatau, M. K., L. D. Talley, and P. P. Niiler (2003), The North Atlantic Oscillation, surface
500 current velocities and SST changes in the subpolar North Atlantic, *16*, 2355–2369.
- 501 Gaillard, F., E. autret, V. Thierry, P. Galaup, C. Coatanoan, and T. Loubrieu (2007),
502 Quality control of large Argo datasets, submitted to *J. Atmos. Ocean. Techn.*, in revi-
503 sion.
- 504 Gonzàlez-Pola, C., A. Lavin, and M. Vargas-Yáñez (2005), Intense warming and
505 salinity modification of intermediate water masses in the southeastern corner of
506 the Bay of Biscay for the period 1992-2003, *J. Geophys. Res.*, *110*, C05020, doi:
507 10.1029/2004JC002367.
- 508 Häkkinen, S., and P. B. Rhines (2004), Decline of Subpolar North Atlantic Circulation
509 during the 1990s, *Science*, *304*, 555–559.
- 510 Hanawa, K., and L. D. Talley (2001), Mode waters, in *Ocean Circulation and Climate*,
511 edited by G. Siedler, J. Church, and J. Gould, pp. 373–386.
- 512 Hátún, H., A. B. Sandø, H. Drange, B. Hansen, and H. Valdimarsson (2005), Influence
513 of the Atlantic Subpolar Gyre on the Thermohaline Circulation, *Science*, *309*, 1841,
514 doi:10.1126/science.1114777.

- 515 Holliday, N. P. (2003), Air-Sea interaction and circulations changes in the northeast At-
516 lantic, *J. Geophys. Res.*, *108*, 3259, doi:10.1029/2002JC001344.
- 517 Hurrell, J. W. (1995), Decadal trends in the North Atlantic Oscillation: regional temper-
518 atures and precipitation, *Science*, *269*, 676–679.
- 519 ICES (2006), Report of the Working Group on Oceanic Hydrography (WGOH), 19-22
520 April 2006, Galway, Ireland, *ICES CM 2006/OCC:08*, 156 pp.
- 521 Johnson, G. C., and N. Gruber (2007), Decadal water mass variations along 20°W in the
522 northeastern Atlantic Ocean, *Prog. Ocean.*, *73*, doi:10.1016/j.pocean.2006.03.022.
- 523 Josey, S. A., and R. Marsh (2005), Surface freshwater flux variability and recent freshening
524 of the North Atlantic in the eastern subpolar gyre, *J. Geophys. Res.*, *110*, CO5008, doi:
525 10.1029/2004JC002521.
- 526 Kwon, Y.-O., and S. C. Riser (2004), North Atlantic Subtropical Mode Water: A his-
527 tory of ocean-atmosphere interaction 1961-2000, *Geophys. Res. Lett.*, *31*, L19307, doi:
528 10.1029/2004GL021116.
- 529 Levitus, S., J. Antonov, and T. Boyer (2005), Warming of the world ocean, 1955-2003,
530 *Geophys. Res. Lett.*, *32*, L02604, doi:10.1029/2004GL021592.
- 531 Lherminier, P., H. Mercier, C. Gourcuff, M. Alvarez, S. Bacon, and C. Kermabon (2007),
532 Transports across the 2002 Greenland-Portugal Ovide section and comparison with
533 1997, *J. Geophys. Res.*, *112*, C07003, doi:10.1029/2006JC003716.
- 534 McCartney, M. S., and L. D. Talley (1982), The Subpolar Mode Water of the North
535 Atlantic Ocean, *J. Phys. Oceanogr.*, *12*, 1169–1188.
- 536 Myers, P. G., N. Kulan, and M. H. Ribergaard (2007), Irminger Water variability in the
537 West Greenland Current, *Geophys. Res. Lett.*, *34*, L17601, doi:10.1029/2007GL030419.

- 538 Pedchenko, A. P. (2005), The role of interannual environmental variations in the geo-
539 graphic range of spawning and feeding concentrations of redfish *Sebastes mentella* in
540 the Irminger Sea, *ICES J. Mar. Sc.*, *62*, 1501–1510.
- 541 Peng, G., E. P. Chassignet, Y.-O. Kwon, and S. C. Riser (2006), Investigation of variability
542 of the North Atlantic Subtropical Mode Water using profiling float data and numerical
543 model output, *Ocean Model.*, *13*, 65–86.
- 544 Perez-Brunius, P., T. Rossby, and D. R. Watts (2004), Transformation of the warm
545 waters of the North-Atlantic from a geostrophic streamfunction perspective, *J. Phys.*
546 *Oceanogr.*, *34*, 2238–2256.
- 547 Pollard, R. T., J. F. Read, N. P. Holliday, and H. Leach (2004), Water masses and
548 circulation pathways through the Iceland Basin during Vivaldi 1996, *J. Geophys. Res.*,
549 *109*, doi:10.1029/2003JC002067.
- 550 Read, J. F. (2001), CONVEX-91: water masses and circulation of the Northeast Atlantic
551 subpolar gyre, *Prog. Ocean.*, *48*, 461–510.
- 552 Reverdin, G., P. P. Niiler, and H. Valdimarsson (2003), North Atlantic Ocean surface
553 currents, *J. Geophys. Res.*, *108*, 3002, doi:10.1029/2001JC001020.
- 554 Reynolds, R., N. Rayner, T. Smith, D. Stokes, and W. Wang (2002), An improved in situ
555 and satellite sst analysis for climate, *15*, 1609–1625.
- 556 Schiermeier, Q. (2007), Artefacts in ocean data hide rising temperatures, *Nature*, *447*,
557 8–9, doi:10.1038/447008a.
- 558 Talley, L. (1999), Mode waters in the subpolar North Atlantic in historical data and during
559 the WOCE period, *Woce News.*, *37*, 3–6.

- 560 Treguier, A. M., S. Theetten, E. Chassignet, T. Penduff, R. Smith, L. Talley, J. O.
561 Beismann, and C. Boening (2005), The North Atlantic subpolar gyre in four high-
562 resolution models, *J. Phys. Oceanogr.*, *35*, 757–774.
- 563 van Aken, H. M., and G. Becker (1996), Hydrography and trough-flow in the northeastern
564 North-Atlantic ocean: The Nansen project, *Prog. Ocean.*, *38*, 297–346.
- 565 Visbeck, M., E. P. Chassignet, R. G. Curry, T. L. Delworth, R. R. Dickson, and K. Krah-
566 mann (2003), The Oceans’s response to North Atlantic Oscillation Variability, in *The*
567 *North Atlantic Oscillation: Climatic Significance and Environmental Impact*, *Geophys.*
568 *Monogr.*, vol. 134, edited by A. G. Union, pp. 113–146.
- 569 Wong, A. P. S., G. C. Johnson, and W. B. Owens (2003), Delayed-mode calibration of
570 autonomous CTD profiling float salinity data by θ -S climatology, *J. Atmos. Ocean.*
571 *Techn.*, *20*, 308–318.
- 572 Yashayaev, I. (2007), Hydrographic changes in the Labrador Sea, 1960-2005, *Prog. Ocean.*,
573 *73*, 242–276, doi:10.16/j.pocean.2007.04.015.

Name	Date	Ship R/V	PI	Reference
NANSEN-90	07/1990	<i>Tyro</i>	Van Aken	<i>van Aken and Becker</i> [1996]
A01E-91	09/1991	<i>Meteor</i>	Meincke	<i>Bersch</i> [1995]
CONVEX-91	08-09/1991	<i>Darwin</i>	Gould	<i>Read</i> [2001]
A01E-92	09/1992	<i>Valdivia</i>	Sy	<i>Bersch et al.</i> [1999]
A01E-95	05-06/1995	<i>Valdivia</i>	Bersch	<i>Bersch et al.</i> [1999]
A01E-96	08-09/1996	<i>Valdivia</i>	Bersch	<i>Bersch et al.</i> [1999]
A01E-97	08-09/1997	<i>Meteor</i>	Sy	<i>Bersch</i> [2002]
OVIDE-02	06-07/2002	<i>Thalassa</i>	Mercier	<i>Lherminier et al.</i> [2007]
OVIDE-04	06-07/2004	<i>Thalassa</i>	Huck	
OVIDE-06	05-06/2006	<i>M. S. Merian</i>	Lherminier	

Table 1. High-quality hydrographic sections used in the analysis.

Figure 1. Bathymetry of the North-Atlantic. The isobaths 0, 200, 1000 and 2000 m are displayed. Points show the positions of the data (Section 2 and Tab. 1): (red) ship-based CTD measurements, (blue) Argo data. Only summer data (June-September) are displayed. Interannual variability of the SPMW properties are estimated in the box delimited by 57.5-59.5°N and 31.5-28.5°W.

Figure 2. (a) NAO Index [*Hurrell*, 1995]. (b, c, d) Anomaly over the period 1950-2006 of air sea fluxes averaged in the eastern subpolar gyre (52-63°N, 33-10°W), derived from the NCEP/NCAR reanalysis 1. (b) Winter (DJFM) total heat flux. A negative heat flux corresponds to a heat loss for the ocean. (c) Winter momentum flux. (d) Annual P-E (Precipitation minus Evaporation). Gray thin lines are winter mean or annual mean values and black thick lines are filtered values using a lanczos filter with a cut-off frequency of 5 years. Averaged values over the entire series are given in each panel.

Figure 3. Potential density along the OVIDE and the A01E sections from Greenland to 25°W in the center part of the Iceland Basin. (a) 2004 OVIDE section. (b) 1992 A01E section. RR indicates the Reykjanes Ridge.

Figure 4. Same as Fig. 3 but for the salinity.

Figure 5. Example of profiles containing Reykjanes Ridge Mode Water. The data were collected in 2002 at 58.41°N– 30.10°W (plain lines) and in 1995 at 58.32°N– 29.94°W (dashed lines). (a) Potential density (black lines) and potential vorticity (gray lines). (b) Potential temperature (black lines) and salinity (gray lines). The mode water core is identified by dots (2002) and squares (1995) and the corresponding properties are indicated in each panels.

Figure 6. Properties of the Subpolar Mode Water in the North Atlantic deduced from hydrographic stations and Argo profiles collected over the period 2001-2006 from beginning of June trough end of September. (a) Potential density. The Reykjanes Box (57.5-59.5°N, 31.5-28.5°W) is indicated. (b) Potential temperature. (c) Salinity.

Figure 7. Time evolution in the Reykjanes Box (57.5-59.5°N, 31.5-28.5°W) of the core properties of the SPMW from 1990 to 2006 estimated from ship-based hydrographic profiles (crosses) and from profiling floats (circles). Black squares are the median properties for each year. The vertical black line represents two times the standard deviation estimated as indicated in the text. Amplitude of the gray bars are proportionnal to the NAO index. (a) Potential density. (b) Potential temperature. The February Reynolds SST averaged in the Reykjanes Box (plain line) and in two boxes located along the Reykjanes Ridge (dashed line: 59.5-61.5°N, 30-26.5°W ; dotted line: 61.5-63.5°N, 25.5-22.5°W) are compared to the temperature of the mode water core. (c) Salinity. (d) Pressure.

Figure 8. Same as Fig. 7 but for the thickness of the mode water layer.

Figure 9. Climatological outcropping position in February of the 27.45 (light gray), 27.5 (dark gray) and 27.55 kg m^{-3} (black) isopycnals deduced from Hydrobase 2 (http://www.whoi.edu/science/PO/hydrobase/HB2_home.htm). The Reykjanes Box and two additional boxes (59.5-61.5°N, 30-26.5°W and 61.5-63.5°N, 25.5-22.5°W), in which we average the February Reynolds SST, are displayed.

Figure 10. February SST from Reynolds' SST monthly fields [*Reynolds et al.*, 2002] : (a) Position of the 7°C isotherm in 1992 (dark blue), 1995 (blue), 1996 (cyan), 1997 (green), 2000 (yellow), 2002 (orange), 2004 (red) and 2006 (brown). (b) February SST in 2004. The thick blue and cyan lines are the 7 and 8°C isotherms, respectively. The contour interval is 0.5°C.

Figure 11. Mean streamfunction for the subpolar North Atlantic from subsurface floats on the 27.5 kg m^{-3} isopycnal. Arrowheads show the direction of flow along contours. The color bar give volume transport for a one-meter-thick layer. The 24,000 and 26,000 $\text{m}^2 \text{s}^{-1}$ contours have been added (dashed). NAC, NWC and IC stand for North Atlantic current, Northwest Corner and Ivinger Current, respectively. See *Bower et al.* [2002] for more details. Copyright Nature.

Figure 12. Oceanic heat content anomaly in the Reykjanes Box (squares) compared to changes in heat content due to atmospheric flux variations only (thick line). In both cases, the 1990-2006 mean is removed.




Figure 1. Bathymetry of the North-Atlantic. The isobaths 0, 200, 1000 and 2000 m are displayed. Points show the positions of the data (Section 2 and Tab. 1): (red) ship-based CTD measurements, (blue) Argo data. Only summer data (June-September) are displayed. Interannual variability of the SPMW properties are estimated in the box delimited by 57.5-59.5°N and 31.5-28.5°W.




Figure 2. (a) NAO Index [Hurrell, 1995]. (b, c, d) Anomaly over the period 1950–2006 of air sea fluxes averaged in the eastern subpolar gyre ($52\text{--}63^\circ\text{N}$, $33\text{--}10^\circ\text{W}$), derived from the NCEP/NCAR reanalysis 1. (b) Winter (DJFM) total heat flux. A negative heat flux corresponds to a heat loss for the ocean. (c) Winter momentum flux. (d) Annual P-E (Precipitation minus Evaporation). Gray thin lines are winter mean or annual mean values and black thick lines are filtered values using a lanczos filter with a cut-off frequency of 5 years. Averaged values over the entire series are given in each panel.




Figure 3. Potential density along the OVIDE and the A01E sections from Greenland to 25°W in the center part of the Iceland Basin. (a) 2004 OVIDE section. (b) 1992 A01E section. RR indicates the Reykjanes Ridge.




Figure 4. Same as Fig. 3 but for the salinity.




Figure 5. Example of profiles containing Reykjanes Ridge Mode Water. The data were collected in 2002 at 58.41°N– 30.10°W (plain lines) and in 1995 at 58.32°N– 29.94°W (dashed lines). (a) Potential density (black lines) and potential vorticity (gray lines). (b) Potential temperature (black lines) and salinity (gray lines). The mode water core is identified by dots (2002) and squares (1995) and the corresponding properties are indicated in each panels.




Figure 6. Properties of the Subpolar Mode Water in the North Atlantic deduced from hydrographic stations and Argo profiles collected over the period 2001-2006 from beginning of June through end of September. (a) Potential density. The Reykjanes Box (57.5-59.5°N, 31.5-28.5°W) is indicated. (b) Potential temperature. (c) Salinity.




Figure 7. Time evolution in the Reykjanes Box (57.5-59.5°N, 31.5-28.5°W) of the core properties of the SPMW from 1990 to 2006 estimated from ship-based hydrographic profiles (crosses) and from profiling floats (circles). Black squares are the median properties for each year. The vertical black line represents two times the standard deviation estimated as indicated in the text. Amplitude of the gray bars are proportionnal to the NAO index. (a) Potential density. (b) Potential temperature. The February Reynolds SST averaged in the Reykjanes Box (plain line) and in two boxes located along the Reykjanes Ridge (dashed line: 59.5-61.5°N, 30-26.5°W ; dotted line: 61.5-63.5°N, 25.5-22.5°W) are compared to the temperature of the mode water core. (c) Salinity. (d) Pressure.




Figure 8. Same as Fig. 7 but for the thickness of the mode water layer.




Figure 9. Climatological outcropping position in February of the 27.45 (light gray), 27.5 (dark gray) and 27.55 kg m⁻³ (black) isopycnals deduced from Hydrobase 2 (http://www.whoi.edu/science/PO/hydrobase/HB2_home.htm). The Reykjanes Box and two additional boxes (59.5-61.5°N, 30-26.5°W and 61.5-63.5°N, 25.5-22.5°W), in which we average the February Reynolds SST, are displayed.




Figure 10. February SST from Reynolds' SST monthly fields [*Reynolds et al.*, 2002] : (a) Position of the 7°C isotherm in 1992 (dark blue), 1995 (blue), 1996 (cyan), 1997 (green), 2000 (yellow), 2002 (orange), 2004 (red) and 2006 (brown). (b) February SST in 2004. The thick blue and cyan lines are the 7 and 8°C isotherms, respectively. The contour interval is 0.5°C.




Figure 11. Mean streamfunction for the subpolar North Atlantic from subsurface floats on the 27.5 kg m^{-3} isopycnal. Arrowheads show the direction of flow along contours. The color bar give volume transport for a one-meter-thick layer. The $24,000$ and $26,000 \text{ m}^2 \text{ s}^{-1}$ contours have been added (dashed). NAC, NWC and IC stand for North Atlantic current, Northwest Corner and Iminger Current, respectively. See *Bower et al.* [2002] for more details. Copyright Nature.




Figure 12. Oceanic heat content anomaly in the Reykjanes Box (squares) compared to changes in heat content due to atmospheric flux variations only (thick line). In both cases, the 1990-2006 mean is removed.